

Formulas for the approximation of the complete Elliptic Integrals

Nikos Bagis

Department of Informatics
Aristotle University Thessaloniki Greece.
e-mail: nikosbagis@hotmail.gr

Abstract

In this article we give evaluations of the two complete elliptic integrals K and E in the form of Ramanujans type- π formulas. The result is a formula for $\Gamma(1/4)^2 \pi^{-3/2}$ with accuracy about 120 digits per term.

keywords: elliptic functions; singular modulus; Ramanujan; Legendre functions; evaluations; constants

1 Elliptic singular moduli

It is known that (see [1],[3])

$$K(x) = \int_0^{\pi/2} \frac{d\theta}{\sqrt{1-x^2 \sin^2(\theta)}} = \frac{\pi}{2} {}_2F_1\left(\frac{1}{2}, \frac{1}{2}; 1; x^2\right) \quad (1)$$

is the complete elliptic integral of the first kind. The function k_r is called elliptic singular moduli and defined from the equation

$$\frac{K\left(\sqrt{1-k_r^2}\right)}{K(k_r)} = \sqrt{r} \quad (2)$$

Also it is known that if $r \in \mathbf{Q}_+^*$, the k_r are algebraic numbers.

For $r \in \mathbf{N}$ we set $K[r] = K(k_r)$. $K[r]$ could be expressed in terms of products of Γ functions, algebraic numbers and powers of π . In time it became obvious that the best way to express the numbers $K[r]$ most concisely was to use the function

$$b(p) = \frac{\Gamma^2(p)}{\Gamma(2p)} \sqrt{\tan(p\pi)} \quad (3)$$

It is also known that if $N = n^2 m$, where n and m are positive integers then

$$K[n^2 m] = M_n(m) K[m], \quad (4)$$

where $M_n(m)$ is algebraic. The following formulas for some $M_n(m)$ are known.

$$M_2(m) = \frac{1 + k'_m}{2} \quad (5)$$

$$27M_3^4(m) - 18M_3^2(m) - 8(1 - 2k_m^2)M_3(m) - 1 = 0 \quad (6)$$

$$(5M_5(m) - 1)^5(1 - M_5(m)) = 256k_m^2(1 - k_m^2)M_5(m) \quad (7)$$

These formulas for finding $K[4r]$, $K[9r]$ and $K[25r]$ depend only on knowing k_r . Also we consider the complete elliptic integral of the second kind, which is

$$E(x) = \frac{\pi}{2} {}_2F_1\left(\frac{-1}{2}, \frac{1}{2}; 1; x^2\right) \quad (8)$$

and related with $K(x)$ from the relation

$$E(k_r) = \frac{K(k_r)}{\sqrt{r}} \left(\frac{\pi}{3K(k_r)^2} - a(r) \right) + K(k_r). \quad (9)$$

The function $a(r)$ is called elliptic alpha function (see [4]).

We will use the elliptic functions theory to evaluate values of $K(k_r)$ and $E(k_r)$ in high precision using Ramanujan's type- π formulas, but now the constant will be not π but

$$\frac{1}{\pi} b \left(\frac{1}{4} \right) = \frac{\Gamma\left(\frac{1}{4}\right)^2}{\pi^{3/2}} \quad (10)$$

the precision of the application formula, which is our more interesting result in this paper is an about 120 digits per term.

Our methods consists Legendre functions, and we not use the function $a(r)$.

2 Legendre polynomials and the formula

The Legendre P function is defined by

$$P_\nu^\mu(z) = \frac{1}{\Gamma(1-\nu)} \left(\frac{z+1}{1-z} \right)^{\nu/2} {}_2F_1\left(-\mu, \mu+1; 1-\nu; \frac{1-z}{2}\right) \quad (11)$$

Set

$$\phi(z) = {}_2F_1(-\mu, \mu+1; 1-\nu; z) = \left(\frac{z}{1-z} \right)^{\nu/2} \Gamma(1-\nu) P_\nu^\mu(1-2z)$$

Then derivating ϕ we have

$$\begin{aligned} \phi'(z) &= \frac{1}{2(1-z)z} \left(\frac{z}{1-z} \right)^{\nu/2} \Gamma(1-\nu) \times \\ &\times [(-1-\mu+\nu+2(1+\mu)z) P_\nu^\mu(1-2z) + (1+\mu-\nu) P_\nu^{1+\mu}(1-2z)] \end{aligned} \quad (12)$$

If we assume that

$$\sum_{n=0}^{\infty} \frac{(-\mu)_n (1+\mu)_n}{(1-\nu)_n} \frac{z^n}{n!} (\alpha n + \beta) = g \quad (13)$$

then

$$\beta\phi(z) + \alpha z\phi'(z) = g$$

From (11),(12) and (13) we have

Theorem 1.

If

$$\alpha = \frac{2(-1+z)}{-1-\mu+\nu+2z+2\mu z} \quad (14)$$

then

$$\sum_{n=0}^{\infty} \frac{(-\mu)_n (1+\mu)_n}{(1-\nu)_n n!} z^n (\alpha n + 1) = \frac{(-1-\mu+\nu) \left(\frac{z}{1-z}\right)^{\nu/2} \Gamma(1-\nu) P_{\nu}^{1+\mu}(1-2z)}{-1-\mu+\nu+2(\mu+1)z} \quad (15)$$

It is known (see [1]), that

$$P_0^{(-1/2)}(1-2z) = {}_2F_1\left(\frac{1}{2}, \frac{1}{2}; 1; z\right) \quad (16)$$

hence if we set $\mu = -3/2$ and $\nu = 0$, then we have

Proposition 1.

$$\sum_{n=0}^{\infty} \frac{\left(\frac{3}{2}\right)_n \left(\frac{-1}{2}\right)_n}{(n!)^2} (k_r)^{2n} [-4(1-k_r^2)n + 1 - 2k_r^2] = \frac{2K(k_r)}{\pi} = 2\vartheta_3^2(q) \quad (17)$$

where $k'_r = \sqrt{1-k_r^2}$, $q = e^{-\pi\sqrt{r}}$.

The result of the Proposition 1 is not trivial since the ϑ_3 -function can be evaluated from the identity

$$\vartheta_3(q) = \sum_{n=-\infty}^{\infty} q^{n^2} \quad (18)$$

in which the two constants e and π involved.

We know that

$$k_{4r} = \frac{1-k'_r}{1+k'_r} \quad (19)$$

Hence

$$K[16r] = \frac{1+k'_{4r}}{2} K[4r] = \frac{1+k'_{4r}}{2} \frac{1+k'_r}{2} K[r]$$

But

$$k'_{4r} = \sqrt{1-k_{4r}^2} = \sqrt{1 - \left(\frac{1-k'_r}{1+k'_r}\right)^2} = \frac{\sqrt{(1+k'_r)^2 - (1-k'_r)^2}}{1+k'_r} =$$

or

$$k'_{4r} = \frac{2\sqrt{k'_r}}{1 + k'_r} \quad (20)$$

Hence

$$K[16r] = \frac{1 + k'_r + 2\sqrt{k'_r}}{4} K[r]$$

or

$$K[16r] = \left(\frac{1 + \sqrt{k'_r}}{2} \right)^2 K[r] \quad (21)$$

Setting $r \rightarrow 4r$ we get

$$K[64r] = \left(\frac{1 + \sqrt{k'_{4r}}}{2} \right)^2 K[4r]$$

or

Lemma.

If $r > 0$, then

$$K[64r] = \frac{\left(\sqrt{1 + k'_r} + \sqrt{2\sqrt{k'_r}} \right)^2}{8} K[r] \quad (22)$$

3 Applications

Set

$$p = 2 + 216 \cdot 5^{1/4} - 96 \cdot 5^{3/4} \quad (23)$$

then

$$k_{100} = \frac{2 - \sqrt{p}}{2 + \sqrt{p}} \text{ and } k'_{100} = \frac{2\sqrt{2}p^{1/4}}{2 + \sqrt{p}} \quad (24)$$

From the duplication formula is

$$k_{400} = \left(\frac{\sqrt{2} - p^{1/4}}{\sqrt{2} + p^{1/4}} \right)^2 \text{ and } k'_{400} = \frac{2^{7/3} p^{1/8} \sqrt{2 + p^{1/2}}}{(\sqrt{2} + p^{1/4})^2}$$

$$k_{1600} = \frac{(\sqrt{2} + p^{1/4})^2 - 2 \cdot 2^{3/4} p^{1/8} \sqrt{2 + \sqrt{p}}}{(\sqrt{2} + p^{1/4})^2 + 2 \cdot 2^{3/4} p^{1/8} \sqrt{2 + \sqrt{p}}}$$

$k_{6400} = w =$

$$= \frac{2 - 2 \cdot 2^{5/8} (2 + \sqrt{p})^{1/4} \sqrt{2\sqrt{2} + 4p^{1/4} + \sqrt{2}\sqrt{p}p^{1/16}} + 2 \cdot 2^{3/4} \sqrt{2 + \sqrt{p}}p^{1/8} + 2\sqrt{2}p^{1/4} + \sqrt{p}}{2 + 2 \cdot 2^{5/8} (2 + \sqrt{p})^{1/4} \sqrt{2 \cdot \sqrt{2} + 4p^{1/4} + \sqrt{2}\sqrt{p}p^{1/16}} + 2 \cdot 2^{3/4} \sqrt{2 + \sqrt{p}}p^{1/8} + 2\sqrt{2}p^{1/4} + \sqrt{p}}$$

Also from (22) we have

$$K[6400] = \frac{1}{8} \left(\sqrt{1 + \frac{2\sqrt{2}p^{1/4}}{2 + \sqrt{p}}} + 2^{7/8} \left(\frac{p^{1/4}}{2 + \sqrt{p}} \right)^{1/4} \right)^2 K[100]$$

But it is known that

$$K[100] = \frac{4 + 2\sqrt{5} + \sqrt{2}(3 + 2 \cdot 5^{1/4})}{80} b\left(\frac{1}{4}\right)$$

hence we get an about 120 digits per term formula for $\frac{1}{\pi}b(1/4)$:

$$\begin{aligned} & \frac{1}{8} [4 + 2\sqrt{5} + \sqrt{2}(3 + 2 \cdot 5^{1/4})]^{-1} \left[\sqrt{1 + \frac{2\sqrt{2}p^{1/4}}{2 + \sqrt{p}}} + 2^{7/8} \left(\frac{p^{1/4}}{2 + \sqrt{p}} \right)^{1/4} \right]^{-2} \times \\ & \times \sum_{n=0}^{\infty} \frac{\left(\frac{3}{2}\right)_n \left(\frac{-1}{2}\right)_n}{(n!)^2} (w)^{2n} [-2(1 - w^2)n - w^2 + 1/2] = \frac{\Gamma\left(\frac{1}{4}\right)}{\pi^{3/2}} \quad : (a) \end{aligned}$$

The evaluation of $E(k_r)/\pi$ follows if we use the formula

$$P_{1/2}(1 - 2z) = \frac{2}{\pi} [2E(z) - K(z)], \quad (25)$$

Then one can arrive with the same method as in Proposition 1, to

Proposition 2.

$$\frac{4E(k_r)}{\pi} = \frac{2K(k_r)}{\pi} + \sum_{n=0}^{\infty} \frac{\left(\frac{1}{2}\right)_n^2}{(n!)^2} (k_r)^{2n} [4(1 - k_r^2)n + 1 - 2k_r^2] \quad (26)$$

References

- [1]: M.Abramowitz and I.A.Stegun, 'Handbook of Mathematical Functions'. Dover Publications
- [2]: B.C.Berndt, 'Ramanujan's Notebooks Part II'. Springer Verlag, New York (1989)
- [3]: B.C.Berndt, 'Ramanujan's Notebooks Part III'. Springer Verlag, New York (1991)
- [4]: J.M. Borwein and P.B. Borwein, 'Pi and the AGM'. John Wiley and Sons, Inc. New York, Chichester, Brisbane, Toronto, Singapore (1987).
- [5]: I.S. Gradshteyn and I.M. Ryzhik, 'Table of Integrals, Series and Products'. Academic Press (1980).
- [6]: E.T.Whittaker and G.N.Watson, 'A course on Modern Analysis'. Cambridge U.P. (1927)
- [7]: I.J.Zucker, 'The summation of series of hyperbolic functions'. SIAM J. Math. Ana.10.192(1979)
- [8]: Bruce C. Berndt and Heng Huat Chan, 'Eisenstein Series and Approximations to Pi'. Page stored in the Web.
- [9]: D. Broadhurst, 'Solutions by radicals at Singular Values k_N from New Class Invariants for $N \equiv 3 \pmod{8}$ '. arXiv:0807.2976v3(math-phy).